

## ADIPOSE MESENCHYMAL STEM CELLS AND “REGENERATIVE ADIPOSE TISSUE GRAFT” (LIPOGEMS™) FOR MUSCULOSKELETAL REGENERATION

C. TREMOLADA<sup>1</sup>, G. BELTRAMI<sup>1</sup>, A. MAGRI<sup>1</sup>, F. BIANCHI<sup>2</sup>, C. VENTURA<sup>2,3</sup>,  
C. DI VITO<sup>4</sup>, R. CAMPANELLA<sup>5</sup>, S.E. NAVONE<sup>6</sup>, G. MARFIA<sup>6</sup> and A.I. CAPLAN<sup>7</sup>

<sup>1</sup>Image Institute, Milan, Italy; <sup>2</sup>Cardiovascular Department, University of Bologna, Italy;  
<sup>3</sup>Laboratory of Molecular Biology and Stem Cell Engineering, National Institute of Biostructures and Biosystems (NIBB), Rome, Italy; <sup>4</sup>Department of Medical Biotechnology and Translational Medicine, LITA-Segrate, University of Milan, Italy; <sup>5</sup>Neurosurgical Unit, San Carlo Borromeo Hospital, Milan, Italy; <sup>6</sup>Laboratory of Experimental Neurosurgery and Cell Therapy, Neurosurgery Unit, Fondazione IRCCS CA' Granda Ospedale Policlinico Milano, Italy; <sup>7</sup>Skeletal Research Center, Department of Biology, CaseWestern Reserve University, Cleveland, USA

Regenerative medicine is a high-potential sector of strategic developments in medicine and health industry. The perspective to cure diseases up to now relied on medical treatments of long duration and limited effectiveness, and the possibility to avoid organ transplantation renders regenerative medicine attractive. In recent years, basic and translation research held great hope for this new field with significant progress in the modulation of stem cell commitment *in vitro* and providing protocols for targeted clinical applications. In line with this approach, mesenchymal stem cells (MSCs) have been introduced as potential therapeutic tools to correct the breakdown of musculoskeletal disorders. MSCs are able to secrete a large number of trophic factors capable of repairing the recipient tissue through angiogenic, anti-apoptotic and anti-fibrotic mechanisms. In this context, adipose tissue is emerging as a clinically relevant and easy to harvest source of multipotent progenitors to develop regenerative therapies. The present review focuses on the clinical application of MSCs, and in particular of adipose-derived stem cells, in the musculoskeletal disorders and on the current scientific challenges. In this perspective, we discuss future developments of an innovative system (Lipogems) for musculoskeletal regeneration, yielding a non-expanded and ready-to-use microfractured fat tissue product harbouring MSCs and pericytes within a preserved stromal vascular niche. The Lipogems system may also pave the way for future off-the-shelf and large-scale approaches for reconstructive procedures and regenerative medicine.

Musculoskeletal diseases are common conditions, including more than 150 different diseases. According to the World Health Organisation, these disorders are the most common cause of severe long-term pain and physical disability (1). With aging of the population, the incidence of musculoskeletal disease is rising and will be a significant socio-

economic burden on society. The complex nature of these conditions generally means that treatment options are limited to managing symptoms, rather than prevention and cure (2). Therefore, there is an urgent need to develop new and effective therapeutic approaches for these age-related

*Key words: musculoskeletal disorders, mesenchymal stem cells, stromal-vascular niche, lipogems*

Corresponding author: Carlo Tremolada, MD,  
President Lipogems International, srl,  
Director Istituto Image,  
v.le Bianca Maria, 24,  
20129 Milan, Italy  
Tel.: +39 3356963371 Fax: +39 02 3651834  
E-mail: carlo.tremolada@gmail.com

disabling pathologies. In this context, great strides have been made in regenerative therapies thanks to stem cell technology. More in detail, recent research has extensively shown the potential of mesenchymal stem cells (MSCs), a class of multipotent stromal stem cells, for reparative/regenerative medicine even in musculoskeletal system (3). Indeed, MSCs isolated from various tissues can differentiate into relevant cell types, thus representing an attractive option for cell-based therapy (4). Moreover, MSCs hold a great promise in tissue restoration thanks to their ability to secrete a wide range of growth factors, which have trophic effects on surrounding host cells, stimulating reparative responses (5). Furthermore, MSCs injected in pathological tissues contribute to favour physiological process by acting as reservoirs of repair cells or immunomodulatory sentinel to reduce inflammation (6). Several early-stage clinical trials are testing the delivery of MSCs in musculoskeletal disease, such as tendon injury, knee osteoarthritis, rheumatoid arthritis. However, despite the excellent potential of MSCs in regenerative medicine, many challenge must be overcome before they can be clinically used (7). In particular, it is essential to standardize protocols of isolation, expansion, and transplantation and to better understand MSC biology. The present review focuses on the clinical application of MSCs in the musculoskeletal disorders, and on the current scientific challenges, in which the difficulty of *ex vivo* expansion and the complexity of the current Good Manufacturing Practice (cGMP) requirements for expanded cells prompt the development of novel approaches in the autologous use of MSCs. In this perspective, we discuss future developments of an innovative system (Lipogems) for musculoskeletal regeneration, yielding a non-expanded and ready-to-use microfractured fat tissue product harbouring MSCs and pericytes within a preserved stromal vascular niche.

## MESENCHYMAL STEM CELLS

### *Biology*

Even if the exact location and function in the tissue of origin is not fully understood, mesenchymal stem cells are defined as multipotent and self-renewable cells with the ability *in vitro* to adhere

to plastic and to differentiate into multiple lineages, including osteogenic, chondrogenic and adipogenic ones (8). Indeed, the International Society for Cellular Therapy define MSCs as a heterogeneous population of progenitor cells expressing a pattern of characteristic, but not specific, surface markers, including CD73, CD90, and CD105, but lacking the expression of hematopoietic markers CD34, CD45, CD14 or CD11, CD79a or CD19, and HLA class II (4). MSCs can be extracted from several body districts, including the adipose and synovial tissues, peripheral blood, skeletal muscle, umbilical cord blood, placenta, and bone marrow. However, an optimal source of MSCs in tissue regeneration and repair has not been yet identified. Indeed, MSCs deriving from different sources have similar, but not equal ability to differentiate into a specific kind of specialized cells. Moreover, the number of cells obtained may depend on the donor age and comorbid conditions.

### *Secretome and stromal-vascular niche*

The participation of MSCs in tissue regeneration has been largely investigated according to the notion that these cells can themselves differentiate into some cell types, including bone, cartilage, muscle, adipocytes, stroma, fibroblasts and endothelial cells (9-11). Recent studies suggest that MSCs could participate in tissue repair, not only differentiating into cells of the target issue, but also releasing several factors, contributing to restorative processes, including angiogenic ones (11-13). Indeed, the secreted trophic factors, participate to tissue rescue through pro-angiogenic and anti-fibrotic mechanisms (14-20), anti-inflammatory and immunomodulatory properties (21-29), anti-apoptotic and antimicrobial characteristics (30-38). Recent studies show a direct correlation between the occurrence of MSCs and the blood vessel density in stromal vascularized tissues (14,39,40). The niche is the morpho-functional unit where stem cells live and reproduce themselves. It is a particular kind of tissue within each tissue, in which a huge network of messages is fashioned through the product of the overall paracrine activity of the embedded cells, the so-called "secretome" (41,42). The regenerative potency of MSCs depends mostly on their ability to afford a timely modulation in the composition of the secretome (43). Further

understanding of the molecular pathways involved in growth factor production will be very helpful to develop better strategies for MSC-based therapies. In this new vision, the preservation of the niche is fundamental to consider MSCs as a patient-specific “molecular biology laboratory” adapting over time to the environmental cues released by the injured tissues (44).

#### *Immunomodulatory effects*

One of the most interesting characteristic of MSCs is their inherent immunomodulatory properties. When transplanted *in vivo*, MSCs do not elicit an immune response, allowing them to be used in allogenic stem cell therapy. In particular, they produce anti-inflammatory cytokines and suppress the proliferation, differentiation and function of immune cells *in vitro* (45). To take full advantage of the unique properties of MSCs for tissue engineering applications, a critical issue will be gaining further insights on the mechanisms controlling their self-renewal and differentiation, which will potentially result in the chance to modulate the behaviour of these cells for therapeutic purposes. Several characteristics of MSCs are purported to impart immune privilege, thereby allowing MSCs to avoid immune rejection in certain situations, which may facilitate the clinical use of allogenic MSCs. MSCs do not express class II Major Histocompatibility Complex (MHC) or costimulator molecules and express low levels of class I MHC (46). One of the first evidence of MSCs role in modulating immune reactions shows that activated MSCs inhibit T-cell expansion in mixed lymphocyte reactions (47). Moreover, MSCs influence the immune system through the secretion of a variety of soluble factors including indoleamine 2,3-dioxygenase (48), nitric oxide (49), transforming growth factor beta (TGF- $\beta$ ), prostaglandin E2 (PGE2) (50), and tumor necrosis factor stimulated gene- Protein (TSG-6) (51). Therefore, in addition to the direct regenerative effect of transplanted MSCs, these cells could interfere with microenvironment to stimulate physiological regeneration. However the precise mechanism of MSC action is not fully elucidate. A deeper comprehension of MSC biology both *in vitro* and *in vivo* could lead to the development of trustworthy strategies for clinical application of MSCs.

#### ADIPOSE MESENCHYMAL STEM CELLS

Adipose tissue is a connective tissue derived from embryonic mesoderm, consisting of a heterogeneous population of cells like adipocytes, preadipocytes, smooth muscle cells, endothelial cells, mast cells, fibroblasts immune cells. About the 10% of the adipocyte population is annually renewed (52). From adipose tissue manipulation it is possible to isolate the so called stromal vascular fraction (SVF), containing, among others, mesenchymal stem cells (53). SVF is easy harvested with minimal donor site morbidity commonly through lipoaspiration followed by *in vitro* cell isolation (54). The efficiency of SVF isolation is strictly related to donor general condition, such as age and obesity (55). In addition, compared to bone marrow, 1 g of adipose tissue contains 500 times more pluripotent cells than 1 g of bone marrow aspirate (54,56). Moreover, besides showing phenotypic and transcriptional profiles similar to that of the other MSCs, ADSCs present some peculiar characteristics. In particular, ADSCs express CD34 glycoprotein (57), stromal markers (CD13, CD29, CD44, CD63, CD73, CD90, and CD166) and endothelial cell markers (CD31, CD144, VEGFR2, and von Willebrand factor) (58). Moreover, ADSCs have a higher yield upon isolation and a greater proliferative rate in culture when compare with MSCs isolated from other sources (59,60). Of note, it has been demonstrated that these cells, besides the multiple mesodermal potential, are also able to differentiate in cells of ectodermal and endodermal origin, considering them as pluripotent stem cells (54,61). Therefore, adipose tissue represents a promising and clinically relevant source of multipotent progenitors to develop regenerative therapies. However, some challenges must be overcome before their use in tissue engineering (7). One of the most relevant problem in the use of ADSCs is their expansion for several weeks *in vitro* prior implantation, that can modify their pluripotent potential and precluding their use in emergency circumstances (62). Moreover, emerging evidence indicates that culturing ADSCs the risk of infection, immunogenicity, genetic instability, and tumorigenicity could increase (63,64). Therefore, there is an urgent need to assess the safety of

ADSC implantation in humans, and to identify novel protocols for ADSC isolation, with minimal manipulation.

#### APPLICATION OF ADSCS FOR MUSCULOSKELETAL REGENERATION

The potential applications of MSCs in regenerative medicine, and, in particular, in musculoskeletal regeneration, show encouraging results for possible clinical applications. However, only few clinical studies use MSC-based approaches for such purposes, due to medical, research, and regulatory reasons (65). However, due to the lack of specific MSC markers, little is known on the *in vivo* ability of MSCs to differentiate. Indeed, different *in vivo* studies show that MSCs could participate to regeneration in various injury models, however, any evidence of clonal expansion, the true differentiation, and regenerative potential does not exist. Therefore, more studies are necessary to elucidate the mechanisms and biological properties of MSCs in determining their therapeutic efficacy in various diseases. Here we summarize the current application of MSCs for musculoskeletal diseases.

##### *Cartilage repair*

Cartilage repair represents a clinical challenge due to the intrinsic limited ability to repair itself. The current treatment of cartilage defects includes intra-articular administration of hyaluronic acid, treatment with platelet-rich-plasma, that contains bioactive proteins such as chemokines, cytokines and growth factors, and bone marrow stimulation, that include subchondral drilling, abrasion, and microfracture. Up to now, the only approved cellular-based therapy for cartilage restoration is based on the autologous chondrocyte implantation after *in vitro* expansion (66). Despite the promising initial results, the limited expansion of chondrocytes *ex vivo*, and donor site morbidity limit the beneficial effects of this technique (66,67). Alternative cellular therapies focused on progenitor cell populations bone marrow stem cells, that once transplanted ameliorated pain and movement ability (68,69). Recently, ADSCs are emerging as a less invasive source of progenitors that can be differentiated into chondrocytes *in vitro* in a 3-dimensional environment (70), in presence of

growth factors (TGF- $\beta$  superfamily) (71) and then implanted to restore cartilage tissue (72). Moreover, it has been observed that also uninduced ADSCs were able to fully restore cartilage in ear auricle defects (73) and in patello patellofemoral joints (74). These promising data suggest, therefore, that a minimal *ex vivo* manipulation of ADSCs, thanks to the intrinsic ability of these cells to adapt to their environment *in vivo*, could allow the development of an easy and effective clinical treatment to restore cartilage defects.

##### *Bone regeneration*

Bone graft is extensively used in a wide range of orthopaedic, plastic oral, dental, and neurocranial surgical procedures (75). However, the use of autograft bone is often associated to morbidity and scarce regenerative properties. Indeed the harvest of bone is associated to pain, haematoma formation, infections, and fractures (76). Moreover, aging reduce the availability of bone marrow stem cells (77). Therefore, research efforts in the field of orthopedics have been directed to ameliorate treatments for reconstructing bony defects, including traumas, tumors, infections, aseptic loosening or nonunions. These investigations have led to the development of improved treatments with strong osteogenic potency, as the recombinant bone morphogenetic proteins (BMPs). BMPs induce and promote critical steps of mostly endochondral bone formation and have been approved for clinical use in spinal fusion, non-unions and severely compromised long bone fractures. However, BMP technology has some therapeutic limitations when the microenvironment is compromised with poor or no vascularization. Therefore, to improve the therapeutic efficacy of this method, BMP administration has been successfully applied together with biomaterials and stem cells (78). In this regard, ADSCs, thanks to their osteogenic potential, have proven as good candidates both *in vitro* and *in vivo* (79,80). Moreover, growing evidence indicate that ADSCs alone, in absence of exogenous growth factors, are able to restore bone defects. In particular, it has been shown that ADSC transplantation could represent a good therapeutic option to treat craniofacial defects (81-83). Surprising results were obtained in the repair of the calvarium, that is unable to ossify after the first

two years of life. ADSCs, administered in presence of milled autologous cancellous bone and fibrin glue, or seeded in  $\beta$ -tricalcium phosphate granules, were able to completely repair calvarial defect, resulting in new bone formation and near complete ossification of the preoperative defect (81,83). However, the use of multiple concomitant treatments limits the comprehension of the therapeutic effect of ADSCs. Indeed, implanted cells could enhance bone regeneration through direct differentiation into mature osteoblasts and by paracrine effects, releasing osteogenic and vasculogenic molecules and growth factors, which facilitate migration and differentiation of resident precursors (84). Nevertheless the promising results obtained with ADSCs in bony defects restorations suggest that cell therapy represents an attractive alternative to traditional treatments like core decompression, osteotomy and total joint replacement, and provide a relatively simple method of autologous bony reconstruction with little donor site morbidity.

#### *Arthritides*

Arthritides, such as rheumatoid arthritis or osteoarthritis, are systemic and disabling autoimmune pathologies characterized by chronic joint inflammation and bone erosion. The crucial process underlying arthritides initiation is the abnormal activation of dendritic cells, T cells, B cells, macrophages, and neutrophils (85). The current treatments for these diseases are mainly addressed to manage symptoms and inflammation. However, a treatment that solves definitively the arthritides does not exist. Some studies focused their attention on cell-based approaches, taking into account that is a systemic defect, differently from focal articular defects, and the underlying disease process. In this context, MSCs, as previously mentioned, hold an immunoregulatory capacity, and elicit immunosuppressive effects, that can be used for several autoimmune diseases, including arthritides. *In vitro* studies on MSCs derived from umbilical cord demonstrated an anti-proliferative effect on synoviocytes, key players in inflammation and joint destruction in rheumatoid arthritis. This effect was mediated by the release of IL-10, 1-methyl-DLtryptophan, TGF- $\beta$ 1, and by the down-regulation of mixed metalloproteinase 9 (86). These results are

in line with the *in vivo* studies on an arthritis rodent model, in which ADSC systemic injection reduced the levels of inflammatory cytokines and chemokines and the ratios of Th1/Th17 cells. Moreover, ADSC transplantation was able to increase the production of the anti-inflammatory IL-10 in lymph nodes and joints, and de novo generation of antigen-specific Treg cells (87,88). Similar results were obtained in the same mouse model after the intraperitoneal infusion of umbilical cord-derived MSCs: arthritis severity was reduced, the levels of proinflammatory cytokines and chemokines were decreased, and IL-10 levels were increased (86). Therefore, taking together these data suggest that ADSCs could ameliorate Arthritis pathogenesis decreasing the production of inflammatory cytokines and activating Treg cells. However, contradictory data was reported in adjuvant-induced and spontaneous arthritis model in which MSCs were effective only when injected before disease onset (89). Therefore, the anti-inflammatory function of stem cells may be effective in preventing or delaying arthritis if delivered at early stages of the disease process. More compelling investigations are needed to improve the specific targeting and the retention of MSCs at the cartilage surface in order to maximize their potential effect.

#### *Tendon*

Tendon injuries are often associated with significant dysfunction and disability, due to the limited self-repair capacity and propensity for scar formation of tendinous tissue. Despite the improvements in conventional treatment, such as transplant, clinical outcomes in tendon treatment are still variable. Moreover the use of allograft can lead to immune response and rejection (90), and the use of autografts is related to donor-site morbidity (91). Therefore, new strategies have been devised, such as tissue engineering techniques. Considering the fact that fibroblasts are involved in tendon healing by producing collagen, in an initial pilot study in humans, *ex vivo* expanded autologous dermal fibroblasts have been successfully used for refractory lateral elbow epicondylitis (92). Recently, with the progression in MSC characteristic identification, tissue engineering relying on these cells has been proposed to enhance tendon healing. In a rat experimental model of patellar tendon window defects, the injection of bone marrow stromal cells in a liquid fibrin matrix

stimulated histological, ultrastructural, molecular, and biomechanical parameters of patellar tendon healing, whereas injection of fibroblasts in fibrin matrix had only minor effects (93). Others *in vivo* studies showed that bone marrow MSCs were effective on tendon-bone healing. In a model of rat Achille's tendon damage, in which the enthesis (bone-tendon junction) was destroyed, the effect of chondrocytes or bone marrow MSCs were compared with not treated rats. Noteworthy, the bone marrow MSCs group showed an enthesis most similar to the premorbid state (94). Similar results were obtained in rabbits undergoing anterior cruciate ligament repair (95). These studies support the application of cell-based therapies for the regeneration of tendon tissues. However, these strategies have been investigated only in pre-clinical studies and the role of stem cells needs to be confirmed.

#### THE CHALLENGE OFFERED BY LIPOGEMS™-DERIVED ADSCs

Lipogems (PCT/IB2011/052204), represent a new completely closed tool to harvest, wash, process, and reinject human (or animal) lipoaspirates. Briefly, the surgical procedure consists in two steps: the infiltration step, in which adrenalin, in a saline solution, and very diluted lidocaine are injected to induce vasoconstriction and local anesthesia, facilitating the subsequent lipoaspiration; the aspiration step, in which a standard liposuction technique is performed. Afterwards, lipoaspirate is processed, by mild mechanical forces, passing floating adipose clusters through different reduction filters 59. Lipogems product is, therefore, a fat tissue derivative minimally manipulated that can be readily injected in an autologous fashion. The fact that Lipogems product is composed of completely normal pericytes (although beginning to detach from the vessels as in NORMAL response to trauma injury from the washing and cluster reduction process) and no cell expansion is done before injection is an extremely safe issue that make this technique just as a standard fat grafting but with less concern regarding the presence of oil and blood contaminations and with smaller size tissue clusters which makes the whole procedure less traumatic.

This device ameliorates the classical Coleman

lipofilling technique (96), eliminating oil and blood residues lipoaspirate and reducing the size of the clusters of adipose tissue (97). These technical improvements have important biologic implications. In islet transplantation for diabetes treatment, the density of loading transplanted tissue and the size of clusters are important predictors of cell transplant outcome, in terms of inflammation, vascularization, and, therefore, in post-transplant engraftment (98). These promising results are probably due to the gentle fat tissue harvesting technique used in Lipogems. Differently from the lipoaspirate, which causes an imbalance in the cytoarchitectonics, Lipogems technique is not traumatic for cells and preserves vascular/stromal architecture with a high percentages of mature and a low amount of hematopoietic-like elements.

In an effort to characterized MSCs resident in fat tissue derivative, it has been observed that Lipogems-derived cells and lipoaspirate cells show a similar tissue architecture. Indeed, any differences in vimentin and fatty acid binding protein 4 distribution, as markers of adipose tissue, and in Ki-67 positivity, as marker of proliferation, were observed (99). Moreover, Bianchi and coworkers demonstrated that the *in vitro* culture of Lipogems derivatives allows the isolation of a population of ADSCs with a high degree of purity, than Lipoaspirate. These cells fulfilled the definition of mesenchymal stem cells, being able to differentiate mesoderm-type of cells, including osteogenic, chondrogenic, and adipogenic ones, like lipoaspirate cells (59). Nevertheless, intriguingly, Lipogems procedure was able to promote the expression of different antigens and to favour the multipotent potential with respect to lipoaspirate cells. In this regard, Carelli and colleagues demonstrated that Lipogems-ADSCs show a higher expression of self-renewal antigens, including OCT4, SOX2, NANOG, and of neural phenotype genes, such as  $\beta$ -tubulin III, NEUROD1, PAX6, and SOX3 (99). Moreover, Lipogems-derived ADSCs, in response to pro-vasculogenic molecules, show a more pronounced expression of angiogenic genes, such as VEGF, KDR, and HGF (59). Furthermore, the treatment with a radio electric asymmetric conveyed field, was able to promote the transcription of genes involved in the commitment towards cardiac, vascular, neuronal,

and skeletal muscle lineages, and of stemness related genes, including Nanog, Sox-2, and Oct-4 (100). The possibility to obtain a final cell product containing viable adipocyte, preadipocyte, and stem cells, eliminating problems related to enzymatic digestion and other manipulations (8), exhibits a great appeal not only for its application in plastic and reconstructive medicine, but also in research, and regenerative medicine. Differently from lipoaspirate, the Lipogems product preserves the stromal vascular fraction also after cryopreservation, bypassing the difficulty of ex vivo expansion and the complexity of current Good Manufacturing Practice requirements for expanded cells. Of interest, the “earlier” cryopreservation of adipose tissue-derived stem cells represents an attractive challenge to treat different degenerative age-related pathologies, as younger adipose tissue progenitors and stem cells have a higher regenerative, tissue remodelling, and therapeutic potential (97,101,102). Based on its properties, the Lipogems product may pave the way to novel approaches and paradigms in the rescue of diseased tissues, within the context of both a personalized and a large-scale regenerative medicine (8).

### CONCLUSIONS

Understanding the dynamics that regulate MSCs homeostasis, especially their anti-inflammatory effect and immunomodulatory capacity, has led to challenge a number of consolidated beliefs on their therapeutic mechanisms. Today, EMA (European Medicines Agency) and FDA (Food and Drug Administration), which had in the past significantly different views in conceiving stem cell biology, efficacy and storage conditions, have developed very similar rules. Nevertheless, these rules are constantly changing, and this is the reason why both Regulatory Agencies refers to “current Good Manufacturing Practice (cGMP)” as the set of guidelines and regulations that should dynamically ensure the currently best available standard for manufacturing medicinal products, including stem cells. However, the emergence of regenerative medicine raises new questions about the best ways to maintain quality. Regenerative medicine therapies involve new kinds of medical products, such as lab-grown or genetically

modified cells. A therapy that uses living cells cannot be standardized in the same way as a conventional pill, and this makes sometimes very difficult to make statements and considerations that may remain valid for more than a few months (94,95). In this complex scenario, the challenge offered by Lipogems represents an attractive technique to obtain minimally manipulated fat tissue product that retains the intact microenvironment in which MSCs live, being amenable for the use in different clinical settings. Moreover, the multipotency of Lipogems-derived MSCs has been shown to be optimized by cell exposure or physical energy providing future off-the-shelf and large scale approaches for reconstructive procedures and regenerative medicine strategy.

**DISCLOSURES.** Prof. C. Tremolada is owner of patents and president of Lipogems International srl.

Prof. C. Ventura is a member of Lipogems International srl.

All other authors have nothing to disclose and declare no potential conflicts of interest with respect to the research, and publication of this article.

### REFERENCES

1. WHO Scientific Group on the Burden of Musculoskeletal Conditions at the Start of the New Millennium. The burden of musculoskeletal conditions at the start of the new millennium. World Health Organ Tech Rep Ser 2003; 919:i-x, 1-218.
2. Brooks P. The burden of musculoskeletal disease - a global perspective. Clin Rheumatol 2006; 25:778-81.
3. Noth U, Rackwitz L, Steinert AF et al. Cell delivery therapeutics for musculoskeletal regeneration. Adv Drug Deliv Rev 2010; 62:765-83.
4. Dominici M, Le Blanc K, Mueller I, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. Cytotherapy 2006; 8:315-7.
5. Caplan AI, Dennis JE. Mesenchymal stem cells as trophic mediators. J Cell Biochem 2006; 98:1076-84.
6. English K. Mechanisms of mesenchymal stromal

- cell immunomodulation. *Immunol Cell Biol* 2013; 91:19-26.
7. Murray IR, West CC, Hardy WR et al. Natural history of mesenchymal stem cells, from vessel walls to culture vessels. *Cell Mol Life Sci* 2014; 71:1353-74.
  8. Murphy MB, Moncivais K, Caplan AI. Mesenchymal stem cells: environmentally responsive therapeutics for regenerative medicine. *Exp Mol Med* 2013; 45:e54.
  9. Prockop DJ. Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* 1997; 276:71-4.
  10. Weissman IL. Translating stem and progenitor cell biology to the clinic: barriers and opportunities. *Science* 2000; 287:1442-6.
  11. Phinney DG, Prockop DJ. Concise review: mesenchymal stem/multipotent stromal cells: the state of transdifferentiation and modes of tissue repair--current views. *Stem Cells* 2007; 25:2896-902.
  12. Chen L, Tredget EE, Wu PYG, Wu Y. Paracrine factors of mesenchymal stem cells recruit macrophages and endothelial lineage cells and enhance wound healing. *PLoS One* 2008; 3:e1886.
  13. Chamberlain G, Fox J, Ashton B, Middleton J. Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells* 2007; 25:2739-49.
  14. Caplan AI. Review: mesenchymal stem cells: cell-based reconstructive therapy in orthopedics. *Tissue Eng* 2005; 11:1198-211.
  15. Petrie Aronin CE, Tuan RS. Therapeutic potential of the immunomodulatory activities of adult mesenchymal stem cells. *Birth Defects Res C Embryo Today* 2010; 90:67-74.
  16. Kolf CM, Cho E, Tuan RS. Mesenchymal stromal cells. *Biology of adult mesenchymal stem cells: Regulation of niche, self-renewal and differentiation. Arthritis Res Ther* 2007; 9:204.
  17. Haynesworth SE, Baber MA, Caplan AI. Cytokine expression by human marrow-derived mesenchymal progenitor cells in vitro: effects of dex- amethasone and IL-1 alpha. *J Cell Physiol* 1996; 166:585-592.
  18. Holgate ST, Davies DE, Lackie PM et al. Epithelial-mesenchymal interactions in the pathogenesis of asthma. *J Allergy Clin Immunol* 2000; 105:193-204.
  19. Doorn J, van de Peppel J, van Leeuwen JPTM et al. Pro-osteogenic trophic effects by PKA activation in human mesenchymal stromal cells. *Biomaterials* 2011; 32: 6089-6098.
  20. Caplan AI, Bruder SP. Mesenchymal stem cells: building blocks for molecular medicine in the 21st century. *Trends Mol Med* 2001; 7:259-64.
  21. Aggarwal S, Pittenger MF. Human mesenchymal stem cells modulate allogeneic immune cell responses. *Transplantation* 2005; 105:1815-22.
  22. Iyer S, Rojas M. Anti-inflammatory effects of mesenchymal stem cells: novel concept for future therapies. *Expert Opin Biol Ther* 2008; 8:569-82.
  23. Uccelli A, Moretta L, Pistoia V. Mesenchymal stem cells in health and disease. *Nat Rev Immunol* 2008; 8:726-36.
  24. Weiss DJ, Bertoncello I, Borok Z et al. Stem cells and cell therapies in lung biology and lung diseases. *Proc Am Thorac Soc* 2011; 8:223-72.
  25. Yagi H, Soto-Gutierrez A, Kitagawa Y. Bone marrow mesenchymal stromal cells attenuate organ injury induced by LPS and burn. *Cell Transplant* 2010; 19: 823-30.
  26. Guan X-J, Song L, Han F-F et al. Mesenchymal stem cells protect cigarette smoke-damaged lung and pulmonary function partly via VEGF-VEGF receptors. *J Cell Biochem* 2013; 114:323-35.
  27. Mantovani A, Sica A, Sozzani S et al. The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* 2004; 25:677-86.
  28. Mantovani A, Sica A, Allavena P et al. Tumor-associated macrophages and the related myeloid-derived suppressor cells as a paradigm of the diversity of macrophage activation. *Hum Immunol* 2009; 70:325-30.
  29. Grivennikov SI, Greten FR, Karin M. Immunity, inflammation, and cancer. *Cell* 2010; 140:883-99.
  30. Cselenyák A, Pankotai E, Horváth EM et al. Mesenchymal stem cells rescue cardiomyoblasts from cell death in an in vitro ischemia model via direct cell-to-cell connections. *BMC Cell Biol* 2010; 11:29.
  31. Li N, Sarojini H, An J, Wang E. Prosaposin in the secretome of marrow stroma-derived neural progenitor cells protects neural cells from apoptotic

- death. *J Neurochem* 2010; 112:1527-38.
32. Kim SY, Lee J-H, Kim HJ et al. Mesenchymal stem cell-conditioned media recovers lung fibroblasts from cigarette smoke-induced damage. *Am J Physiol Lung Cell Mol Physiol* 2012; 302:L891-L908.
  33. Frohm M. The expression of the gene coding for the antibacterial peptide LL-37 is induced in human keratinocytes during inflammatory disorders. *J Biol Chem* 1997; 272:15258-63.
  34. Wu H, Zhang G, Minton JE et al. Regulation of cathelicidin gene expression: induction by lipopolysaccharide, interleukin-6, retinoic acid, and salmonella enterica serovar typhimurium infection. *Infect Immun* 2000; 68:5552-8.
  35. Rogan MP, Geraghty P, Greene CM et al. Antimicrobial proteins and polypeptides in pulmonary innate defence. *Respir Res* 2006; 7:29.
  36. Bonfield TL, Lennon D, Ghosh SK et al. Cell based therapy aides in infection and inflammation resolution in the murine model of cystic fibrosis lung disease. *Stem Cell Discovery* 2013; 03:139-53.
  37. Haniffa MA, Wang X-N, Holtick U et al. Adult human fibroblasts are potent immunoregulatory cells and function- ally equivalent to mesenchymal stem cells. *J Immunol* 2007; 179:1595-604.
  38. Meisel R, Brockers S, Heseler K et al. Human but not murine multipotent mesenchymal stromal cells exhibit broad-spectrum antimicrobial effector function mediated by indoleamine 2,3-dioxygenase. *Leukemia* 2011; 25:648-54.
  39. Da Silva Meirelles L, Sand TT, Harman RJ et al. MSC frequency correlates with blood vessel density in equine adipose tissue. *Tissue Eng Part A* 2009; 15:221-9.
  40. Blocki A, Wang Y, Koch M et al. Not all MSCs can act as pericytes: functional in vitro to distinguish pericytes from other mesenchymal stem cells in angiogenesis. *Stem Cells Dev* 2013; 22:2347-55.
  41. Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell. *Blood Cells* 1978; 4:7-25.
  42. Spradling A, Drummond-Barbosa D, Kai T. Stem cells find their niche. *Nature* 2001; 414:98-104.
  43. Caplan AI. Adult mesenchymal stem cells for tissue engineering versus regenerative medicine. *J Cell Physiol* 2007; 213:341-7.
  - Moore KA, Lemischka IR. Stem Cells and Their Niches. *Science* 2006; 311:1880-5.
  44. Jiang XX, Zhang Y, Liu B et al. Human mesenchymal stem cells inhibit differentiation and function of monocyte-derived dendritic cells. *Blood* 2005; 105:4120-6.
  45. Le Blanc K, Tammik C, Rosendahl K et al. HLA expression and immunologic properties of differentiated and undifferentiated mesenchymal stem cells. *Exp Hematol* 2003; 31:890-6.
  46. Bartholomew A, Sturgeon C, Siatskas M et al. Mesenchymal stem cells suppress lymphocyte proliferation in vitro and prolong skin graft survival in vivo. *Exp Hematol* 2002; 30:42-8.
  48. Krampera M, Cosmi L, Angeli R et al. Role for interferon-gamma in the immunomodulatory activity of human bone marrow mesenchymal stem cells. *Stem Cells* 2006; 24:386-98.
  49. Ren G, Zhang L, Zhao X et al. Mesenchymal stem cell-mediated immunosuppression occurs via concerted action of chemokines and nitric oxide. *Cell Stem Cell* 2008; 2:141-50.
  50. English K, Barry FP, Field-Corbett CP, Mahon BP. IFN-gamma and TNF-alpha differentially regulate immunomodulation by murine mesenchymal stem cells. *Immunol Lett* 2007; 110:91-100.
  51. Lee RH, Pulin AA, Seo MJ et al. Intravenous hMSCs improve myocardial infarction in mice because cells embolized in lung are activated to secrete the anti-inflammatory protein TSG-6. *Cell Stem Cell* 2009; 5:54-63.
  52. Rigamonti A, Brennand K, Lau F, Cowan CA. Rapid cellular turnover in adipose tissue. *PLoS One* 2011; 6:e17637.
  53. Gimble JM, Katz AJ, Bunnell BA. Adipose-derived stem cells for regenerative medicine. *Circ Res* 2007; 100:1249-60.
  54. Zuk PA, Zhu M, Ashjian P et al. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell* 2002; 13:4279-95.
  55. Cawthorn WP, Scheller EL, MacDougald OA. Adipose tissue stem cells meet preadipocyte commitment: going back to the future. *J Lipid Res* 2012; 53:227-46.
  56. di Summa PG, Kalbermatten DF, Pralong E et al. Long-term in vivo regeneration of peripheral nerves

- through bioengineered nerve grafts. *Neuroscience* 2011; 181:278-91.
57. Pittenger MF, Mackay AM, Beck SC et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999; 284:143-7.
  58. Mitchell JB, McIntosh K, Zvonic S et al. Immunophenotype of human adiposederived cells: temporal changes in stromal-associated and stem cell-associated markers. *Stem Cells* 2006; 24:376-85.
  59. Bianchi F, Maioli M, Leonardi E, et al. A new nonenzymatic method and device to obtain a fat tissue derivative highly enriched in pericyte-like elements by mild mechanical forces from human lipoaspirates. *Cell Transplantation* 2013; 22:2063-77.
  60. DeUgarte DA, Alfonso Z, Zuk PA, et al. Differential expression of stem cell mobilization-associated molecules on multi-lineage cells from adipose tissue and bone marrow. *Immunol Lett* 2003; 89:267-70.
  61. Zuk PA. The adipose-derived stem cell: looking back and looking ahead. *Mol Biol Cell* 2010; 21:1783-7.
  62. Ma T. Mesenchymal stem cells: From bench to bedside. *World J Stem Cells* 2010; 2:13-7.
  63. Dahl JA, Duggal S, Coulston N, et al. Genetic and epigenetic instability of human bone marrow mesenchymal stem cells expanded in autologous serum or fetal bovine serum. *Int J Dev Biol* 2008; 52:1033-42.
  64. Torsvik A, Røslund GV, Svendsen A et al. Spontaneous malignant transformation of human mesenchymal stem cells reflects cross-contamination: putting the research field on track - letter. *Cancer Res* 2010; 70:6393-6.
  65. Steinert AF, Rackwitz L, Gilbert F et al. Concise review: the clinical application of mesenchymal stem cells for musculoskeletal regeneration: current status and perspectives. *Stem Cells Transl Med* 2012; 1:237-47.
  66. Brittberg M, Lindahl A, Nilsson A et al. Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation. *N Engl J Med* 1994; 331:889-95.
  67. Diaz-Romero J, Gaillard JP, Grogan SP et al. Immunophenotypic analysis of human articular chondrocytes: changes in surface markers associated with cell expansion in monolayer culture. *J Cell Physiol* 2005; 202:731-42.
  68. Nejadnik H, Hui JH, Feng Choong EP et al. Autologous bone marrow-derived mesenchymal stem cells versus autologous chondrocyte implantation: an observational cohort study. *Am J Sports Med* 2010; 38:1110-6.
  69. Wakitani S, Mitsuoka T, Nakamura N, et al. Autologous bone marrow stromal cell transplantation for repair of full-thickness articular cartilage defects in human patellae: two case reports. *Cell Transplant* 2004; 13:595-600.
  70. Salibian AA1, Widgerow AD, Abrouk M, Evans GR. Stem cells in plastic surgery: a review of current clinical and translational applications. *Arch Plast Surg* 2013; 40:666-75.
  71. Hennig T, Lorenz H, Thiel A et al. Reduced chondrogenic potential of adipose tissue derived stromal cells correlates with an altered TGFbeta receptor and BMP profile and is overcome by BMP-6. *J Cell Physiol* 2007; 211:682-91.
  72. Lin Y, Luo E, Chen X et al. Molecular and cellular characterization during chondrogenic differentiation of adipose tissue-derived stromal cells in vitro and cartilage formation in vivo. *J Cell Mol Med* 2005; 9:929-39.
  73. Bahrani H, Razmkhah M, Ashraf MJ et al. Differentiation of adipose-derived stem cells into ear auricle cartilage in rabbits. *J Laryngol Otol* 2012; 126:770-4.
  74. Zhang HN, Li L, Leng P, et al. Uninduced adipose-derived stem cells repair the defect of full-thickness hyaline cartilage. *Chin J Traumatol* 2009; 12:92-7.
  75. Giannoudis PV, Dinopoulos H, Tsiridis E. Bone substitutes: an update. *Injury*. 2005; 36(S):S20-7.
  76. Finkemeier CG. Bone-grafting and bone-graft substitutes. *J Bone Joint Surg Am* 2002; 84-A:454-64.
  77. Rodríguez JP, Astudillo P, Ríos S, Pino AM. Involvement of adipogenic potential of human bone marrow mesenchymal stem cells (MSCs) in osteoporosis. *Curr Stem Cell Res Ther* 2008; 3:208-18.
  78. Cancedda R, Bianchi G, Derubeis A et al. Cell therapy for bone disease: A review of current status. *Stem Cells* 2003; 21:610-9.
  79. Zuk PA, Zhu M, Mizuno H et al. Multilineage cells

- from human adipose tissue: implications for cell-based therapies. *Tissue Eng* 2001; 7:211-28.
80. Cowan CM, Shi YY, Aalami OO et al. Adipose-derived adult stromal cells heal critical-size mouse calvarial defects. *Nat Biotechnol* 2004; 22:560-7.
  81. Lendeckel S, Jodicke A, Christophis P et al. Autologous stem cells (adipose) and fibrin glue used to treat widespread traumatic calvarial defects: case report. *J Craniomaxillofac Surg* 2004; 32:370-3.
  82. Mesimaki K, Lindroos B, Tornwall J et al. Novel maxillary reconstruction with ectopic bone formation by GMP adipose stem cells. *Int J Oral Maxillofac Surg* 2009; 38:201-9.
  83. Thesleff T, Lehtimaki K, Niskakangas T et al. Cranioplasty with adipose-derived stem cells and biomaterial: a novel method for cranial reconstruction. *Neurosurgery* 2011; 68:1535-40.
  84. Barba M, Cicione C, Bernardini C et al. Adipose-derived mesenchymal cells for bone regeneration: state of the art. *Biomed Res Int* 2013; 2013:416391.
  85. Firestein GS. Evolving concepts of rheumatoid arthritis. *Nature* 2003; 423:356-61.
  86. Liu Y, Mu R, Wang S et al. Therapeutic potential of human umbilical cord mesenchymal stem cells in the treatment of rheumatoid arthritis. *Arthritis Res Ther* 2010; 12:R210.
  87. Gonzalez MA, Gonzalez-Rey E, Rico L et al. Treatment of experimental arthritis by inducing immune tolerance with human adipose-derived mesenchymal stem cells. *Arthritis Rheum* 2009; 60:1006-19.
  88. Zhou B, Yuan J, Zhou Y et al. Administering human adipose-derived mesenchymal stem cells to prevent and treat experimental arthritis. *Clin Immunol* 2011; 141:328-37.
  89. Papadopoulou A, Yiangou M, Athanasiou E et al. Mesenchymal stem cells are conditionally therapeutic in preclinical models of rheumatoid arthritis. *Ann Rheum Dis* 2012; 71:1733-40.
  90. Crossett LS, Sinha RK, Sechriest VF, Rubash HE. Reconstruction of a ruptured patellar tendon with Achilles tendon allograft following total knee arthroplasty. *J Bone Joint Surg Am* 2002; 84:1354-61.
  91. Chiou HM, Chang MC, Lo WH. One-stage reconstruction of skin defect and patellar tendon rupture after total knee arthroplasty. A new technique. *J Arthroplasty* 1997; 12:575-9.
  92. Connell D, Datir A, Alyas F et al. Treatment of lateral epicondylitis using skin-derived tenocyte-like cells. *Br J Sports Med* 2009; 43:293-8.
  93. Hankemeier S, Hurschler C, Zeichen J et al. Bone marrow stromal cells in a liquid fibrin matrix improve the healing process of patellar tendon window defects. *Tissue Eng Part A*. 2009;15:1019-30.
  94. Nourissat G, Diop A, Maurel N et al. Mesenchymal stem cell therapy regenerates the native bone-tendon junction after surgical repair in a degenerative rat model. *PLoS One* 2010; 5:e12248.
  95. Lim JK, Hui J, Li L et al. Enhancement of tendon graft osteointegration using mesenchymal stem cells in a rabbit model of anterior cruciate ligament reconstruction. *Arthroscopy* 2004; 20:899-910.
  96. Allen RJ Jr, Canizares O Jr, Scharf C et al. Grading lipoaspirate: is there an optimal density for fat grafting? *Plast Reconstr Surg* 2013; 131:38-45.
  97. Tremolada C, Palmieri G, Ricordi C. Adipocyte transplantation and stem cells: Plastic surgery meets regenerative medicine. *Cell Transplant* 2010; 19:1217-23.
  98. Tremolada C. *Chirurgia plastica del viso e del collo. Principi generali e guida per il medico estetico.* In: Massirone, ed. *Trattato di Medicina Estetica.* Padova, Italy: Edizioni Piccin-Nuova Libreria 2010; 1855-74.
  99. Carelli S, Messaggio F, Canazza A et al. Characteristics and Properties of Mesenchymal Stem Cells Derived from Micro-fragmented Adipose Tissue. *Cell Transplant* 2014.
  100. Maioli M, Rinaldi S, Santaniello S et al. Radio electric asymmetric conveyed fields and human adipose-derived stem cells obtained with a non-enzymatic method and device: a novel approach to multipotency. *Cell Transplant* 2013; 23(12):1489-500.
  101. Oishi K, Noguchi H, Yukawa H, et al. Cryopreservation of mouse adipose tissue-derived stem/progenitor cells. *Cell Transplant* 2008; 17:35-41.
  102. Alt EU, Senst C, Murthy SN et al. Aging alters tissue resident mesenchymal stem cell properties. *Stem Cell Res* 2012; 8:215-25.